

Supplemental Labeling



EPA Reg. No. 5905-564

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR THIS PRODUCT BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. "Label" as used in this supplemental labeling refers to the label booklet for this product and this supplemental.

KEEP OUT OF REACH OF CHILDREN DANGER – PELIGRO

GENERAL INFORMATION

LATIGO[®] is a post-emergence herbicide for control of a wide spectrum of annual, biennial, and perennial broadleaf weeds in teff production.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions and Conditions of Sale and Warranty are to be followed. This labeling must be in the user's possession during application.

APPLICATION TO TEFF

LATIGO[®] may be applied by ground or air.

For ground applications, apply in 10 to 20 gallons of total broadcast spray mix per acre. For aerial application, apply in 3 to 10 gallons of water per acre.

The Restricted Entry Interval (REI) is 48 hours.

The Pre-harvest Interval (PHI) is 14 days.

Limited to one postemergence application per crop cycle.

Limited to one preharvest application per crop cycle.

Refer to the label for all crop rotation restrictions.

GRASS SEED CROPS

LATIGO[®] application rates and spray volumes will vary with the growth stage and population of broadleaf weeds to be controlled. In general the smaller the weed, the lower use of the recommended rate range will provide satisfactory control.

The larger the weed, the population and environmental conditions will require the higher end of the rate range to achieve the satisfactory control especially for many of the perennial weeds.

To control emerged broadleaf weeds, apply 0.75 – 1.25 pints of **LATIGO**[®] per acre. Apply after the two-leaf stage and prior to the boot stage. Refer to "Plant Response" and "Livestock Feeding Restrictions" under the **GRASS PASTURES** section of the label.

PASTURES, RANGELAND AND GRASS (Hay, Silage)

In newly established Teff, use 0.75 to 1.25 pints of **LATIGO**[®] per acre to control or suppress weeds after planting. Newly seeded areas including small grains grown for pasture or hay, may be injured if rates of **LATIGO**[®] are greater than 1.25 pints per acre are applied. Apply after the two-leaf stage and prior to the boot stage.

Grazing and Feeding Non-Lactating Animals: There is no waiting period between treatment and grazing for non-lactating animals. Do not permit meat animals being finished for slaughter to graze treated fields within 30 days of slaughter.

Grazing and Feeding Lactating Animals: Do not graze lactating dairy animals within 7 days of treatment.

Dry Hay and Silage: Treated grasses may be harvested for dry hay or silage but do not harvest within 37 days of treatment.

SMALL GRAIN CROPS (Grains used for Gluten-Free Flour)

Apply up to 1.25 pints of **LATIGO**[®] per acre to Teff. Apply after the two-leaf stage and prior to the boot stage. If small grains are grown for pasture or hay only, refer to Pastures, Rangeland and Grass (Hay, Silage).

HELENA CHEMICAL COMPANY
225 Schilling Blvd., Suite 300
Collierville, Tennessee 38017
Telephone: 901-761-0050

AD 012612

LATIGO[®] is a registered trademark of Helena Holding Company.
©Helena Holding Company, 2010.